

THERAPEUTIC POTENTIAL OF PLANTS IN WOUND HEALING

Rajinder Raina, A.K. Srivastava, Shahid Prawez and Azad Ahmad

Division of Pharmacology and Toxicology,
F.V.Sc. & A.H., SKUAST-J, R.S. Pura, Jammu

Introduction

Wound is defined simply as the disruption of the cellular and anatomic continuity of a tissue (Bennet 1988). Wound may be produced by physical, chemical, thermal, microbial or immunological insult to the tissue. The process of wound healing consists of integrated cellular and biochemical events leading to reestablishment of structural and functional integrity with regain of strength of injured tissue. The events include coagulation, inflammation, formation of granulation tissue and tissue remodeling (Lynch 1987; Savant & Shah 1998).

Clinically, one often encounters non-healing, under-healing or over healing. Therefore the aim of treating a wound is to either shorten the time required for healing or to minimize the undesired consequences (Myers *et al* 1980). Attention is directed towards discovering an agent, which will accelerate wound healing either when it is progressing normally (Brown *et al.*, 1988a; Mather *et al.*, 1989), or when it is suppressed by various agents like corticosteroids (Ehrlich & Hunt 1968), antineoplastics (Raju & Kulkarni 1986), non steroidal anti-inflammatory agents (Lee 1968 b).

Medical treatment of wound includes administration of drugs either locally (topical) or systemically (oral or parenteral) in an attempt to aid wound repair (Savanth & Shah 1998; Rains & Mann 1988; Moy 1993). The topical agents used include antibiotics and antiseptics (Chulani 1996), desloughing agents (chemical debridement, e.g. Hydrogen peroxide, eusol and collagenase ointment) (Savanth & Mehta 1998), wound healing promoters (eg. Tretinoin, *aloe vera* extract, honey, comfrey, benzoyl peroxide, *chamomilia* extract, dexpanthenol, tetrachlordecaxide solution, clostebol acetate and the experimental cytokines).

Various growth factors like platelet derived growth factor, macrophage derived growth factor, monocyte derived growth factor (Brown *et al.*, 1988a; Mather *et al.*, 1989) etc. are necessary for the initiation and promotion of wound healing. Many substances like tissue extracts (Udupa *et al* 1991; Ramesh *et al.*, 1990), vitamins (Williams & Bissel 1944), minerals (Rao *et al.*, 1988a) and a number of plant products (Dahanukar *et al.*, 2000; Sharma *et al.*, 1990; Udupa *et al.*, 1991); have been reported by various workers, to possess prohealing effects. These approaches are being advocated in wound management as body's repair kit (McAuliffe 1989).

Consequently there exists a need for new agents which may be useful in proper wound management. In this direction a number of herbal products are being investigated.

Herbal products for wound management

Herbal products have been used in wounds management for many years. A few plants/plant products with promising outcome are discussed here.

Aloe vera

Aloe, native to Africa, is also known as “lily of the desert”, the plant of immortality; and the medicinal plant. The name was derived from the alloeh meaning “bitter” because of the bitter liquid found in the leaves. In 1500 B.C Egyptians recorded use of this herbal plant in treating burns, infections and parasites.

Extensive research has shown that the clear gel of it has a dramatic ability to heal wounds, ulcers, burns by putting a protective coating on the affected areas and speeding up the healing process. The plant contains 96% of water and rest is essential oil, aminoacids, minerals, vitamins, enzymes and glycoproteins. Various constituents of *Aloe vera* have been shown to have anti-inflammatory activity as well as to simulate wound healing (Penneys 1981).

There are reports that *Aloe vera* gel applied topically could help heal radiation burns. However, a large modern, placebo controlled study did not find aloe effective in this regard (Williams *et al.*, 1996). Some clinical reports suggest topical *Aloe vera* gel for healing minor burns (Visuthilosol *et al.*, 1995).

Aloe vera, topical application of the gel is harmless and hypersensitive reactions to it are rare. In some severe burns, aloe gel may actually impede healing (Schmidt & Greenspoon 1991).

Lantana camara

Lantana camara Linn (Fam-Verbenaceae) a shrub, native of tropical America and completely naturalized in many parts of India as an ornamental plant. The plant has wound healing, abortifacient, antimalarial and anti-inflammatory properties (Kirtikar & Basu 1994; Chopra *et al.*, 1956; Kurian 1995).

The leaves of the plant are reported to be useful in the treatment of wounds, ulcers, bruises, sores etc. (Brando *et al.*, 1985). In a recent study hydroalcoholic extract and fresh juice of leaves have favored wound contraction (Dash *et al.*, 2001).

The toxicities of the plant include nephrotoxicity, hepatotoxicity, photosensitization, dermatitis, intestinal haemorrhages (Adesina 1982; Sharma *et al.*, 1992). In view of the alarming reports of the toxicity, the use of this plant in whole or any part thereof needs to be carefully regulated until the alarming toxic principles of the plant are properly identified and removed.

St. John's Wort

St. John's wort is a bushy perennial plant with numerous yellow flowers. It is native to many parts of the world including Europe and the United States. It has a 2400 years history of safe and effective usage in many folk and herbal remedies. It is claimed to be useful in mental depression, anxiety, sleep disorders, menstrual cramping, sciatica and arthritis. The blossoms have been used in folk medicine to relieve ulcers, gastritis, diarrhea and nausea.

Externally it is used on cuts as disinfectants and to relieve inflammation and promote healing. The tincture of *Hypericum* spp. given orally has a remarkable effect in lacerated and suppurated wounds, and play an important role in restoring tissue vitality (Fayazuddin 1981).

Further studies have also confirmed *Hypericum* spp. tincture has pro-healing action (Rao *et al.*, 1991), as evidenced by the increase in wound contraction rate and granulation tissue breaking strengths. In addition, epithelization phase was also enhanced.

Tridax procumbens

The plant is a native of tropical America and naturalized in tropical Africa, Asia, Australia and India. Leaf of *Tridax procumbens* mainly contains crude protein 26%, crude fiber 17% soluble carbohydrate 39% calcium oxide 5% (Chadha 1976).

The juice of the leaves of this plant is used by villagers to arrest bleeding from cuts and bruises in animals. It has been shown that its juice accelerates two phases of healing namely epithelization and collagenization; and retards scar formation and granulation (Diwan *et al.*, 1982). From these observations it was postulated that *Tridax procumbens* perhaps have certain components with prohealing and some with antihealing properties.

Tridax procumbens antagonized anti-epithelization and tensile strength depressing effect of dexamethasone (a known healing suppressant agent) without affecting anticontraction and antigranulation action of dexamethasone (Diwan *et al.*, 1983). The effect of various extracts (whole plant extract, aqueous extract, butanol extract and ether fraction) of this plant has been studied in dead space wound model (Udupa *et al.*, 1991).

The authors have reported that whole plant extract has the greatest prohealing activity with increase in tensile strength and lysyl oxidase activity among the various extracts. Aqueous extract was also effective in increasing lysyl oxidase but to a lesser degree than whole plant extract.

Further it has been shown that extract of leaves of this plant also promotes wound healing in both normal and immunocompromised (steroid treated) rats in dead space wound model. The plant increased not only lysyl oxidase but also, protein and nucleic acid content in the granulation tissue, probably as a result of increase in glycosamino glycan content (Udupa *et al.*, 1998).

Chromolaena odorata

Chromolaena odorata was first identified in central America and Vietnam. The aqueous extract and the decoction from leaves of this plant have been used throughout Vietnam for the treatment of soft tissue wounds and burn wounds.

In clinical use it has been noted that aqueous extracts of *Chromolaena odorata* enhances hemostatis (Akah 1990) and stimulates granulation tissue and reepithelization (Lee 1995). The extract also inhibits wound contraction and the inhibition of contraction is reversible. This supports that suggestion that it might be of therapeutic value in minimizing post burn scar contracture and deformities.

Hydnocarpus wightiana

The oil of *Hydnocarpus* spp. has been recognized for several years as anti-leprosy drug and as an anti-parasitic drug in the treatment of guinea worm infestation. There are reports that wounds in leprosy patients and in patients with diabetic ulcers and gangrene healed faster when the oil of *Hydnocarpus* spp. was given orally or administered topically (Manjrekar 1996).

Recently the wound healing effect of oil of *Hydnocarpus* spp. was studied with reference to collagenation and the strength of scar tissue (Oomen *et al.*, 1999). The drug treated group showed a significant increase in strength of scar tissues in the incision wound model and also increased the strength of collagen tissue and hydroxyproline content in the dead space wound model.

Hydnocarpus oil administered orally promoted epithelization, but not wound contraction (Oomen *et al.*, 2000). External application of oil of *Hydnocarpus* spp. and its paste significantly shortened the epithelization period when compared to control group. Further it was suggested that this finding is useful clinically because the oil may act as adjuvant in healing of wounds and ulcer in leprosy patients.

***Helianthus annus* Linn.**

An ornamental annual herb, with erect, rough and hairy stem, common in Indian Gardens in swampy areas. In traditional medicine the plant is used by tribals for inflammation of eyes, sores, dysuria, colic, bite of tigers and bone fractures (Jain & Tarafdar 1970). The seed oil contains high levels of lysine, arginine, aspartic and glutamic acids and tannins (Jain *et al.*, 1974).

The alcoholic extract of whole plant of *H. annus* applied in the form of an ointment on the excised wound of rat led to a significant reduction in total healing period. This has been confirmed by histology where earlier appearance of fibroblasts are seen. Further it is stated that *H. annus* has hastened repair process which is indicated by a high degree of accumulation and an early appearance of the mucopolysaccharide (Deshpande *et al.*, 1965).

Jasminum auriculatum

A small herb found in south India and the western peninsula. The alcohol free defatted extract of *J. auriculatum* leaves has been reported to contain lupeol and jasminol (Deshpande & Upadyaya 1967). *J. auriculatum* has been shown to be beneficial in wound healing. The juice of the leaves in the form of jelly, on local application to a linear uniform excised wound in rats was found to promote wound healing, as assessed by histological, biochemical and contraction rate studied (Deshpande *et al.*, 1965).

Further, it has been reported that the fresh juice of the leaves showed an increased and early gain of the tensile strength in the treated linear wounds in rats. The study indicated that collagenation contributed to improved tensile strength in the early phase of healing (Deshpande & Pathak 1965).

Further studied on musculo-peritoneal wound on the abdomen of rats, with 2.5% leaves extract of the plant injected intramuscularly confirmed the prohealing property of leaves. The effect was more marked in case of skin wounds than in the musculoperitoneal wounds (Deshpande & Pathak 1966a).

Ghee medicated with *J. auriculatum*, on topical application was found to accelerate the healing time of second degree burn wounds in rats by six days. The mucopolysaccharide accumulation was found to be significantly higher in group treated with medicated ghee.

The acceleration of the healing time was approximately 20% with unmedicated ghee and 30% in medicated ghee (Deshpande & Pathak 1966b).

Ginkgo biloba

Ginkgo biloba (*Salisburia adiantifolia*) is also known as maiden hair tree. The genus ginkgo originated 200 million years ago and is considered as a living fossil (Kubitzki K. 1990; Hori *et al.*, 1997). Extracts of leaves have been used therapeutically for centuries (Newall *et al.*, 1996).

The main constituent are flavonoids and terpenes trilactones. *Ginkgo biloba* exhibits a variety of interesting pharmacological activities such as increase in blood fluidity, antioxidant, prohealing,

membrane stabilizing and improvement in cognition (Clostrew 1986; Kleijin & Knipschild 1992; Bairy & Rao 2001). *Ginkgo biloba* has promoted epithelization without altering wound contraction. In case of dead space wounds *Ginkgo biloba* has increased granulation tissue breaking strength without altering granulation tissue mass weight.

However, it did significantly enhance the content of hydroxyl-proline of granulation tissue. Further it is reported that the prohealing action of the *Ginkgo biloba* is due to the presence of flavonoids (Bairy & Rao 2001).

Septilin

A proprietary herbal preparation claimed to be helpful in Gram negative and Gram positive infections (Gadekar *et al.*, 1986; Sharma SK *et al.*, 1986). It consists of *Balsamodendron mukul* (guggul), extracts *Maharasanadi kwath*, extract *Phyllanthus embica*, extract *Tinospora cordifolia*, *Rubia cordifolia*, extract *Glycyrrhiza glabra* and shanka bhasma which are claimed to have wound healing promoting action. Septilin has promoted gain in tensile strength in incision wound model, but at the same time, did not modify the granulation phase of healing (Udupa *et al.*, 1989).

In case of excision wound Septilin promoted epithelization and wound contraction (Nadakarni & Nadakarni 1954). This may be due its effect on migration and mitosis of epithelial cells and promotion of contraction of myo-fibroblasts, the later being now recognized as responsible for wound contraction (Gibbiani *et al.*, 1992).

Centella asiatica

Centella asiatica (Brahmi) also known as "gotu kola", is the main herb in Ayurveda for nervous system, it is used in the repair of nervous tissue from crushing trauma, such as spinal injury, neuromuscular disorders, and to increase general brain function and memory concentration. Used extensively in Asia for the treatment of leprosy, gotu kola heals a host of skin conditions, including wounds, cellulites, varicose vein.

Only the aqueous extract suspension in 5% propylene glycol of *Centella asiatica* as compared to other extracts (viz. alcoholic, petroleum ether, chloroform extract) promoted wound healing in experimentally induced open wounds on topical administration in rats as evidenced by the increase in collagen content and thickness of epithelium (Rao *et al.*, 1996).

Suguna *et al* 1998 also demonstrated that alcoholic extract (oral and topical) of *Centella asiatica* improved the rate of wound healing in rats. It has also been shown that topical administration of the aqueous extract increased cellular proliferation, promoted the collagen synthesis at the wound site as evidenced by the increase in DNA, protein, collagen content of granulation tissue and in tensile strength. The treated wound epithelized faster as compared to control.

Among the various formulations (ointment, cream and gel) of aqueous extract, the process of healing was better with gel formulation (Kumar *et al.*, 1998). It has also been demonstrated that active principles (Triterpens and asiaticoside) promote rapid wound healing (Maquart *et al* 1999; Sukla *et al* 1999).

Cissus quadrangularis

Cissus quadrangularis (Asthisankala) found in hotter parts of India. It is also wide spread in the drier parts of Africa and Arabia. Powdered roots are used for fracture of bone. The effect of the

total extract of the plant on the healing of cortisone treated fracture has been compared with that of anabolic steroid nandrolone. Total extract of the plant, on parenteral administration, was found to neutralized anti-anabolic effect of cortisone in healing fractures.

The stimulatory effect of the total extract was found to be greater than that of anabolic steroid probably due to its vitamin content (Prasad & Udapa 1963). The total extract was found hasten fracture healing by reducing the total convalescence period by 33% in experimental rats and dogs and produced a quick recovery in animals. The up-take of calcium is quick which helps in early completion of calcification process and earlier remodeling phenomenon.

The tensile strength studies showed an early gain, leading to a 90% gain of its normal strength at the end of six weeks in comparison to 60% gain in strength noted in controls (Udapa & Prasad 1964). The plant extract has been found to influence fracture healing indirectly. Its effect on organic and inorganic phase of bone repair in normal and some pathological conditions like alloxan diabetes and in endocrine response during fracture healing suggested that parenteral administration of drug acted directly on the testes and through pituitary to release the androgenic hormone in to blood.

The other evidence of its androgenic property was its ability to produce a positive nitrogen balance, increase in body weight and total weight of testis of animals. A potent anabolic steroid isolated from the plant showed a marked influence on the rate of fracture healing by influencing early ingeneration process of all the connective tissue involved in the healing and quick mineralization of the callus (Satyavathi *et al.*, 1976).

Recently it has been reported that methanolic extract of *Cissus quadrangularis* promoted the healing process of experimentally fractured radius-ulna of dogs as evidenced by radiologically and histopathological examinations. The treated group also exhibited a reduction in serum calcium levels as compared to saline control animals (Deka *et al.*, 1994).

Miscellaneous prohealers from plants

The effects of *Azadirachata indica*, *Ocimum sanctum*, and *Begia odorata* on infected experimental wounds in laboratory animals have been reported. All these plants, notably *Ocimum sanctum* promote healing (Thaker & Anjaria 1986). *Euphorbia nerrifolia* (aqueous extract) when applied topically facilitated the healing of surgically produced cutaneous wounds in guinea pigs. It has increased the gain in tensile strength, DNA content and promoted epithelization (Rasik *et al.*, 1996).

Both crude betal nut extract and its polyphenols promoted healing of incision and dead space wounds (Padmaja *et al.*, 1993). Alcoholic extract of *Indigofera aspalathoids* has anti-inflammatory, wound healing and analgesic effect (Bhaskar *et al.*, 1982).

Fresh leaves of *Kalanchoea integra* showed encouraging results in inflammatory wound healing (Yadav & Yadav 1985). Mango butter which is extracted from the seeds of *Mangifera indica* has a wound healing property. It is used as application for ulcerations, fissures of lips, hands and chapped skin. Anecdotal evidence, some clinical observations, some animal model studies and few randomized clinical trials support the efficacy of honey in managing wounds (Molan 1998; Molan 1999).

Reference:

- Adesina SK. (1982) *Fitoterapia*, 53: 147-162.
- Akah PA. (1990) *Int. J. Crude Drug Res.* 28(4): 253-256.
- Atal CK, Srivastava JB, Wali BK, Shankaran PS. *Indian J. Exp. Biol.* 16: 330-333.
- Bairy KL, Rao CM. (2001) *J. Nat. Remed.* 1: 25-27.
- Bennet RG. (1988). *Fundamentals of cutaneous surgery*. St. Louis: C.V.Mosby; 778.
- Bhaskar EA, Ganga N, Arivudainambi R, Shanthanam G. (1982) *Indian J. Med. Res.* 76: (Suppl) 115.
- Brando M, Boteltho M, Krettli E. (1985) *Cienc. Cult.* 37(7): 1152-1163.
- Brown GL, Curtsinger LT, White M. (1988a). *Annals of Surgery* 208: 788-794.
- Chadha YR. (1976) *Wealth of India*, 10: 292.
- Chopra RN, Nayar SL, Chopra IC. (1956) *Glossary of Indian Medicinal plants.*, Publication and Informational Directorate, CSIR: New Delhi; 149-150.
- Chulani HL. (1996) In: *The law of medical negligence* 1st edn. Radhakrishnan Medical and Educational Trust: Mumbai; 51-83.
- Clostrew. (1986) *Press Med.* 15: 1529-1538.
- Dahanukar SA; Kulkarni Ra, Rege NN. (2000) *Indian J. Pharmacol.* 32: S81-S118.
- Dash GK, Suresh P, Ganapathy S. (2001) *J. Nat. Remed.* 1(2); 105-110.
- Deka DK, Lahon LC, Saikia J, Mukit A. (1994) *Indian J. Pharmacol.* 26: 45-46.
- Deshpande PJ, Pathak SN, Shankaran PS (1965) *Indian J. Med. Res.* 53: 539.
- Deshpande PJ, Pathak SN. (1965) *Surg. J. Delhi*; 1: 273.
- Deshpande PJ, Pathak SN. (1966a) *Med. Surg.* 6: 21.
- Deshpande PJ, Pathak SN. (1966b) *Indian J. Med. Res.* 1(1): 81.
- Deshpande PJ, Shankaran PS, Pathan SN. (1965) *Med. Surg.* 5: 27.
- Deshpande SM, Upadyaya RR. (1967) *Curr Sci.* 36: 233.
- Diwan PV, Tillo LD, Kulkarni DR. (1982) *Indian J. Med. Res.* 75; 460-466.
- Diwan PV, Tillo LD, Kulkarni DR. (1983) *Indian J. Physiol. Pharmacol.* 27(1): 32-36.
- Ehrlich HP, Hunt TK. (1968) *Ann Surg.* 167: 324-326.
- Fayazuddin M (1981) *Faiz*, Homeopathic Publication House: Kakinada; 30.
- Gadekar H, Vijay A, Jyoti, Viba, Komawar, Sonavar SG. (1986) *Probe*, 25: 164.
- Gibbiani G, Hirschel BJ, Ryan GB, Statkov PR, Majno G. (1972) *J. Exp. Med.* 135: 719-734.
- Hori T, Ridge RW, Tuleckew, DeIT. (1997) Guiller JT, Tobe H. Springer-Verlag: Tokyo; 350.
- Jain NC, Shivalkar ND, Rao BY. (1974) *J. Oil Technol. Assoc. India*, 6:60-62.
- Jain SK, Tarafdar CR. (1970) *Medicine plant love of Sautals* (A review of P.O. Boddington's work) *EconBot.* 24: 241.
- Kirtikar Kr, Basu BD. (1994) *Indian Medicinal plants*, Bishen Singh Mahendrapal Singh: Dehradun; 1913-1915.
- Kleijin J, Knipschild P. (1992) *The Lancet* 340: 1136-1139.
- Kubitzki K. (1990) Kramer KV, Green PS. (Eds.) Springer-verlag: Berlin; 284-289.
- Kurian JC. (1995) *Plants that heal*, Owners Oriental Watchman Publishing House: Pune; 190.
- Lee KH, (1968 b) *J. Pharm. Sci.* 57: 1238-1240.

- Lee TT (1995) *the use of Eupolin prepared from Eupatorium to treat soft tissue wounds*. The 5th European Tissue Repair Society Meeting, Padova, Italy.
- Lynch WS. (1987) In: Epstein E, Epstein E. (Eds.) *Skin Surgery*, 6th edn., WB Saunders: Philadelphia; 56-70.
- Manjrekar S. (1996) *10th Annual report of the Samidha Charitable Trust*, p 10.
- Maquart FX, Castang F, Simeon A, Birenbant, Gillery P, Wegrowski Y. (1999) *Eur. J. Dermatol.* 9(4): 289-296.
- Mather MD, Sherman M, Frycakowski A, Jester JV. (1989) *Invest Ophthalmol. Vismal Sci.* 30: 2403-2406.
- McAuliffe. (1989) *Span* April 18-19.
- Molan PC. (1998) *Primary intention* 6(4): 148-158.
- Molan PC. (1999) *J. Wound Care*, 8(8): 423-426.
- Moy LS. (1993) *Detmatol. Clin.* 11: 759-766.
- Myers KA, Marshal RD, Friedin J. (1980). *Principles of Pathology in Surgery*, 1st edn. Blackwell Scientific Publications: London; 58-82.
- Nandakarni KM, Nadakarni AK. (1954) *Indian Materia Medica III rd edn*. Popular book depot, Doota Paperhwar Prakashan Ltd: Bombay; 167-170.
- Newall CA, Anderson La, Philipson JD. (1996) *ginkgo in herbal medicine, a guide for health care professionals*, The Pharmaceutical Press: London; 138-140.
- Oomen ST, Rao CM, Raju CVN. (1999) *Int J. Lepr.* 67(2): 154-158.
- Oomen ST, Rao CM, Raju CVN. (2000) *Int J. Lepr.* 68(1): 69-70.
- Padmaja PN, Bairy KL, Kulkarni DR. (1993) *Fitoterapia* LXV(4): 298-300.
- Penneys NS. (1981) *Acta Derm Stockh*, 62: 59-61.
- Prasad GC, udapa KN. (1963) *Indian J. Med. Res.* 51: 667-671.
- Rains AJH, Mann CV. (1988) InL Rains AJH, Mann CV. (Eds) *Bailey and Love's A short practice of surgery*, 20th edn. English Language Book Society: London; 1-11.
- Raju SS, Kulkarni DR. (1986) *Indian J Pharmacol*, 18: 154-157.
- Ramesh KV, Rao CM, Bairy KL, Kulkarni DR. (1990) *Indian J. Exp. Biol.* 28: 43-45.
- Rao CM, Ashok Kumar, Kulkarni DR. (1988a) *Indian J. Physiol. Pharmacol.* 32(1): 61-66.
- Rao SG, Udapa AL, Udapa SL, Rao PGM, Rao Ganesh, Kulkarni DR. (1991) *fitoterapia* LXII(6): 508-510.
- Rao VG, Shivakumar HG, Parthsarathi G. (1996) *Indian J. Pharmacol.* 28: 249-253.
- Rasik AM, Shukla A, Patnail GK. (1996) *Indian J. Pharmacol.* 28: 107-109.
- Satyavathi GV, Raina MK, Saran M. (1976) *Medicinal Plants of India*, Vol. 1, 1st edn. Cambridge Press: New Delhi; 242.
- Savant SS, Shah RA. (1998) In: Savant SS, Shah RA, Gore D. (Eds.) *Text book and atlas of Dermatology and Cosmetology*, 1st edn. Mumbai: ASCAD; 12-17.
- Savanth SS, Mehta N. (1998) In: Savanth SS, Shah RA, Gore D. (Eds.) *Text book and atlas of dermatosurgery and cosmetology*, 1st edn. ASCAD: Mumbai; 50-61.

- Savanth SS, Shah RA. (1998) In: Savanth SS, Shah RA, Gore D. (Eds.) *Text book and atlas of dermatosurgery and cosmetology*, 1st edn., ASCAD: Mumbai; 1-11.
- Schmidt JM, Greenspoon JS. (1991) *Obstet. Gynecol.* 78: 115-117.
- Sharma OP, Makkar HPS, Dawra RK. (1982) *Xenobiotica*, 12(4): 265-269.
- Sharma SK, Agrawal HC, Dharam Pal, Bikchandani. (1986) *Probe*, 25: 156.
- Sharma SP, Aithal Ks, Srinivasan KK, Udupa AL, Vasanth K, Kulkarni DR. (1990) *Fitoterapia* LXI: 263-266.
- Shukla A, Rasik AM, Dhavan BN. (1999) *Phytother. Res.* 13(1): 50-54.
- Suguna L, Shivakumar P, Chandra kasan G. (1998) *Indian J. Exp. Biol.* 34: 1208-1211.
- Sunil K, Parameshwaraiah S, Shivakumar HG. (1998) *Indian J. Exp. Biol.* 36: 569-572.
- Thaker HM, Anjaria JV. (1986) *Indian J. Pharmacol.* 18: 171-174.
- Udupa KN, Prasad GC. (1964) *Indian J. Med. Res.* 52: 482-485.
- Udupa SL, Rao SG, Kulkarni DR. (1989) *Indian J. Physiol. Pharmacol.* 33:1-5.
- UDupa SL, Shaila HP, Udupa AL, Ramesh KV, Kulkarni DR. (1991) *biochem Arch.* 7: 207-212.
- Udupa SL, Udupa AL, Kulkarni DR. (1991) *Fitoterapia* LXII: 46-150.
- Udupa SL, Udupa AL, Kulkarni DR. (1991) *Planta Med.* 57: 325-327.
- Udupa SL, Udupa AL, Kulkarni DR. (1998) *Fitoterapia* 69: 507-510.
- Visuthilosol V, Chowchuen B, Sukwanarat Y. (1995) *J. Med. Assoc. Thai*, 78: 403-409.
- Williams MS, Berk M, Lopriniz CL. (1996) *Int. J. Rad. Oncol. Boil. Phys.* 36: 345-349.
- Williams RH, Bissel GW. (1944) *Arch. Surg.* 49: 225-230.
- Yadav CL, Yadav CS. (1985) *Ancient Sci. Aife.* 5: 30.